## **CLAIMS**

## What is claimed is:

- 1. An isolated polypeptide molecule comprising residues 26 to 546 of SEQ ID NO:2.
- 2. The isolated polypeptide molecule according to claim 1, wherein the polypeptide molecule comprises residues 26 to 627 of SEQ ID NO:2.
- 3. The isolated polypeptide molecule according to claim 1, wherein the polypeptide molecule comprises residues 1 to 627 of SEQ ID NO:2.
- 4. The isolated polypeptide molecule of claim 1, wherein at least nine contiguous amino acid residues of SEQ ID NO:2 are operably linked via a peptide bond or polypeptide linker to a second polypeptide selected from the group consisting of maltose binding protein, an immunoglobulin constant region, and a polyhistidine tag.
- 5. The isolated polypeptide molecule according to claim 1, wherein the polypeptide comprises a fusion protein wherein polypeptide is conjugated with a compound selected from the group consisting of keyhole limpet hemocyanin, muramyl dipeptide, histidine-tag, beta gal, and palmitic acid.
- 6. An isolated polynucleotide molecule encoding the polypeptide molecule according to claim 1.
- 7. An expression vector comprising the following operably linked elements:
  - a) a transcription promoter;
  - b) a DNA segment encoding the polypeptide according to claim 63 and

- c) a transcription terminator.
- 8. An expression vector of claim 7 wherein the DNA segment further encodes an affinity tag.
- 9. A cultured cell into which has been introduced an expression vector according to claim 7, wherein said cell expresses the polypeptide encoded by the DNA segment.
- 10. A method of producing a polypeptide comprising culturing a cell according to claim 9, whereby said cell expresses the polypeptide encoded by the DNA segment, and recovering the polypeptide.
  - 11. The polypeptide made by the method of claim 10.
- 12. A method of producing an antibody to a polypeptide comprising the following steps:

inoculating an animal with the polypeptide such that the polypeptide elicits an immune response in the animal to produce the antibody; and

isolating the antibody from the animal,

wherein the polypeptide is chosen from,

- a) a polypeptide comprising residues 26 to 546 of SEQ ID NO:2;.
- b) a polypeptide comprising residues 26 to 627 of SEQ ID NO:2; and
- c) a polypeptide comprising residues 1 to 627 of SEQ ID NO:2 and wherein the antibody produced by the method specifically binds to a polypeptide of SEQ ID NO:2.
  - 13. The antibody produced by the method according to claim 12.

14. A method of producing an antibody to a polypeptide comprising the following steps:

inoculating an animal with the polypeptide such that the polypeptide elicits an immune response in the animal to produce the antibody; and

isolating the antibody from the animal,

wherein the polypeptide comprises at least fifteen consecutive amino acids of the amino acid sequence as shown in SEQ ID NO:2,

and wherein the antibody produced by the method specifically binds to a polypeptide of SEQ ID NO:2.

- 15. The antibody produced by the method according to claim 14.
- 16. A method for inhibiting sperm-oocyte fusion comprising contacting the sperm and oocytes with the polypeptide according to claim 1, or a fragment thereof, whereby the polypeptide or fragment inhibits the sperm and oocyte fusion.
- 17. A method for inhibiting sperm-oocyte fusion comprising contacting the sperm and oocytes with the antibody according to claim 12, whereby the polypeptide or fragment inhibits the sperm and oocyte fusion.
- 18. A method for inhibiting sperm-oocyte fusion comprising contacting the sperm and oocytes with the antibody according to claim 14, whereby the polypeptide or fragment inhibits the sperm and oocyte fusion.
- 19. A method for inducing infertility in a mammal, comprising administering to the mammal an contraceptively effective dose of the polypeptide according to claim 1, or a fragment thereof, wherein the polypeptide induces an immune response in the mammal, thereby inducing infertility in the mammal.

- 20. A method for inducing infertility in a mammal, comprising administering to the mammal the antibody according to claim 12, wherein the polypeptide induces an immune response in the mammal, thereby inducing infertility in the mammal.
- 21. A method for inducing infertility in a mammal, comprising administering to the mammal the antibody according to claim 14, wherein the polypeptide induces an immune response in the mammal, thereby inducing infertility in the mammal.
- 22. A composition comprising a contraceptive dose of the polypeptide according to claim 1, or a fragment thereof, and an acceptable carrier, and/or adjuvant.